



Thyroid screening strategies in Fasting, postmeal & random blood samples in prolonged diabetics

¹Dr Arun Tadas, Professor & Head , Biochemistry Department, Indira Gandhi Government Medical College, Nagpur, Maharashtra, India

²Dr Monali Rewatkar, Assistant Professor, Biochemistry Department, Indira Gandhi Government Medical College, Nagpur, Maharashtra, India

³Mr Sanjay Agrawal, statistitian PSM, Indira Gandhi Government Medical College, Nagpur, Maharashtra, India

Corresponding Author: Dr Monali Rewatkar, Assistant Professor, Biochemistry Department, Indira Gandhi Government Medical College, Nagpur, Maharashtra, India

Citation this Article: Dr Arun Tadas, Dr Monali Rewatkar, Mr Sanjay Agrawal, “Thyroid screening strategies in Fasting, postmeal & random blood samples in prolonged diabetics”, IJMSIR- February - 2020, Vol – 5, Issue -1, P. No. 01 – 06.

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

Abstract

Background: Diabetes mellitus (DM) and thyroid dysfunction (TD) are the two most common endocrine disorders in clinical practice . Patients with diabetes mellitus are at an increased risk of developing thyroid diseases.[1] The frequency of thyroid dysfunction in diabetic patients is higher than that of the general population and up to a third of patients with diabetes (DM) ultimately develop thyroid dysfunction. Thyroid dysfunction may impair metabolic control and add to cardiovascular disease risk in prolonged diabetic patients.[2] The objective of this study was to investigate the prevalence of Thyroid dysfunction in patients with diabetes mellitus.

Aims: Our aim is to study Pearsons correlation between thyroid dysfunction in diabetes mellitus and to examine thyroid disease screening strategies in routine diabetes care, in diabetic patients (fasting,postmeal,Random blood samples) attending clinical biochemistry OPD at tertiary care Unit, IGGMC,Nagpur.

Materials & Methods: This study includes about 150 patients attending the clinical Biochemistry OPD, IGGMC, Nagpur. All Diabetic patients (more than 5 yrs) who were attending biochemistry OPD for routine Blood sugar tests were selected for this study. History was taken as per designed proforma and consent form was obtained. We collect 50 Fasting Blood samples ,50 Postmeal & 50 Random blood samples from known Diabetic patients .Collected sample was analysed for Blood sugar as well as thyroid test ie T3,T4 & TSH. Blood sugar test was run on Autoanalyser EM 460 in clinical Biochemistry Laboratory, IGGMC, Nagpur. Thyroid tests was run on Elisa reader & washer Serum values of Thyroid stimulating hormone (TSH), thyroxine (T4) and tri-iodo thyronine (T3) were assayed by ELISA tests and values were compared & correlated with Blood sugar level in Diabetics in all groups Fasting, Postmeal as well as Random Blood samples. The data was analysed & Pearsons Correlation was obtained. Student's T-test was used & . P value <0.05 was considered statistically significant.

Results: Results of our study shows Positive Pearsons correlation between TSH level & Blood sugar in Fasting, Postmeal & Random Blood samples of all Diabetic Patients while there is Negative Pearsons correlation between T3,T4 in all Diabetics. P value is found to be statistically significant in all groups (cases)

Conclusions: We conclude that screening for thyroid disease among patients with diabetes mellitus should be routinely performed considering the prevalence of new cases diagnosed and the classical risk factors such as hypertension and dyslipidemia, arising from an undiagnosed thyroid dysfunction.

Keywords : Blood Sugar level, Fasting, Post meal & Random, Thyroid Profile Test,Prolonged Diabetes.

Introduction

Diabetic patients have a higher prevalence of thyroid disorders compared with the normal population Diabetes Mellitus (DM) and thyroid dysfunction (TD) are the two most common endocrine disorders in clinical practice [3]. The association between DM and TD is widely known, with the first studies published in 1979 [4]. Since then, several studies in different countries were conducted to estimate the prevalence of TD in diabetic patients. The relationship between TD and DM is characterized by a complex interaction of interdependence. Screening of TD, especially the subclinical dysfunction, in patients with DM is justified because most patients can be asymptomatic.[5]

The strong link between diabetes and thyroid diseases encouraged the American Diabetes Association (ADA) to propose that people with diabetes must be checked periodically for thyroid dysfunction [6]. Thyroid disease should be screened annually in diabetic patients to detect asymptomatic thyroid dysfunction [7].

Objectives of the study

- 1) The aim of this study is to investigate the prevalence of TD in patients with diabetes mellitus (DM)
- 2) To study Pearsons correlation between Blood sugar level in prolonged diabetes & thyroid profile
- 3) To study correlation between fasting, postmeal, random blood sugar with thyroid dysfunction in prolonged diabetics

Method of collection of data & selection of subjects

This study include about 150 patients attending the clinical Biochemistry OPD, IGGMC, Nagpur. All Diabetic patients (history of Diabetes more than 5 yrs) who were attending biochemistry OPD for routine Blood sugar tests were selected for this study. History was taken as per designed proforma and consent form was obtained. We collect 50 Fasting Blood samples , 50 Postmeal & 50 Random blood samples from known Diabetic patients Collected sample was analysed for Blood sugar as well as thyroid test ie T3,T4 & TSH.

Serum values of Thyroid stimulating hormone (TSH), thyroxine (T4) and tri-iodo thyronine (T3) were assayed by ELISA tests and values were compared & correlated with Blood sugar level in Diabetics in all groups Fasting,Postmeal as well as Random Blood samples.

Blood sugar test was run on Autoanalyser EM 460 in clinical Biochemistry Laboratory,IGGMC,Nagpur.

Blood sugar level was estimated by GOD-POD kit method.

Normal Range – Fasting (70 to 100 mg%) Postmeal (upto 150 mg%) Random (70 to 120 mg%)

Thyroid tests was run on Elisa reader & washer.

For T3, T4, TSH – Immunoassay Elisa kit method on Elisa Reader & Washer

Normal Range –T3 - 0.52 to 1.85 ng/ml, T4 - 5 to 15 ug/dl, TSH – 0.39 to 6.16 uIU/ml

Analysis was carried on Autoanalyser EM – 460 in clinical Biochemistry lab, IGGMC for Serum cholesterol, Triglycerides, HDL,LDL & VLDL . All estimations was done & their values were compared & correlated. The data was analysed & Pearsons Correlation was obtained. Student's T-test was used & . P value <0.05 was considered statistically significant.

Results

Blood Sugar	Correlation	T3	T4	TSH
Glucose fasting	Pearson Correlation	-0.703	-0.760	0.718
	P value	0.000	0.000	0.000
Glucose Post meal	Pearson Correlation	-0.330	-0.446	0.368
	P value	0.001	0.000	0.000
Glucose Random	Pearson Correlation	-0.396	-0.510	0.426
	P value	0.000	0.000	0.000

All p values are significant

	T3	T4	TSH	glucose(f)	glucose(PM)	glucose®
mean	1.0427	4.554	5.634	103.96	169.53	129.95
S.D	0.340982	2.65403904	4.706416	22.77156	34.24819	26.37851

Discussion

Result of our study shows TSH level (Mean – 5.63) in Diabetic patients in all blood samples ie fasting,postmeal & random is positively correlated with Blood sugar level in Diabetes by Pearsons correlation (Table 1) Similarly Vice versa with T3 ,T4 & Blood sugar level which shows Negative Pearsons Correlation. Our findings similar to findings from Spain published by Díez et al. in 2011 [8], Effects of thyroid hormones on various metabolic processes like diabetes mellitus may worsen diseased condition.. Uncontrolled hyperthyroidism in diabetic patients may trigger hyperglycaemic emergencies while recurrent hypoglycaemic episodes have been reported in diabetic patients with hypothyroidism.[9]

Furthermore, thyroid dysfunction may amplify cardiovascular disease risk in diabetic patients through inter-relationships with dyslipidaemia, insulin resistance and vascular endothelial dysfunction.[10]

Our result of a positive correlation between blood sugar level and TSH is consistent with the results by Hollowell,et al. [11].

In a study by M. F. Celani et al. [12], 89 % of patients had hypothyroidism and 11 % had hyperthyroidism. Hypothyroidism was shown to be more prevalent thyroid disorder in type 2 diabetics in the studies of Strieder, M. F. Prummel et al. [13]. The increased frequency of thyroid dysfunction in diabetic patients and its likely deleterious effects on cardiovascular and metabolic function calls for a systematic approach to thyroid disease screening in diabetes.[14] Routine annual thyroid testing should be targeted at diabetic patients at risk of thyroid dysfunction such as patients with DM.

Conclusion

Thyroid dysfunction is common in diabetic patients and can produce significant metabolic disturbances. The presence of thyroid dysfunction may affect diabetes control. Therefore, regular screening for thyroid abnormalities in all diabetic patients will allow early treatment and prevent further complications as well as improve diabetic control which can be otherwise adversely affected by thyroid dysfunction.

References

1. L. H. Duntas, J. Orgiazzi, and G. Brabant, “The interface between thyroid and diabetes mellitus,” *Clinical Endocrinology*, vol. 75, no. 1, pp. 1–9, 2011. View at Publisher · View at Google Scholar · View at Scopus
2. J. L. Johnson, “Diabetes control in thyroid disease,” *Diabetes Spectrum*, vol. 19, no. 3, pp.

- 148–153, 2006. View at Publisher · View at Google Scholar · View at Scopus
3. D. H. Akbar, M. M. Ahmed, and J. Al-Mughales, “Thyroid dysfunction and thyroid autoimmunity in Saudi type 2 diabetics,” *Acta Diabetologica*, vol. 43, no. 1, pp. 14–18, 2006. View at Publisher · View at Google Scholar · View at Scopus
 4. J. Feely and T. E. Isles, “Screening for thyroid dysfunction in diabetics,” *British Medical Journal*, vol. 1, no. 6179, p. 1678, 1979. View at Google Scholar · View at Scopus
 5. P. Perros, R. J. McCrimmon, G. Shaw, and B. M. Frier, “Frequency of thyroid dysfunction in diabetic patients: value of annual screening,” *Diabetic Medicine*, vol. 12, no. 7, pp. 622–627, 1995. View at Google Scholar · View at Scopus
 6. American Diabetes Association, “Standards of medical care in diabetes—2013,” *Diabetes Care*, vol. 36, no. 1, pp. S11–S66, 2013. View at Publisher · View at Google Scholar
 7. A.-R. M. Radaideh, M. K. Nusier, F. L. Amari et al., “Thyroid dysfunction in patients with type 2 diabetes mellitus in Jordan,” *Saudi Medical Journal*, vol. 25, no. 8, pp. 1046–1050, 2004. View at Google Scholar · View at Scopus
 8. J. J. Díez, P. Sánchez, and P. Iglesias, “Prevalence of thyroid dysfunction in patients with type 2 diabetes,” *Experimental and Clinical Endocrinology and Diabetes*, vol. 119, no. 4, pp. 201–207, 2011. View at Publisher · View at Google Scholar · View at Scopus
 9. R. Kadiyala, R. Peter, and O. E. Okosieme, “Thyroid dysfunction in patients with diabetes: clinical implications and screening strategies,” *International Journal of Clinical Practice*, vol. 64, no. 8, pp. 1130–1139, 2010. View at Publisher · View at Google Scholar · View at Scopus
 10. R. S. Gray, W. J. Irvine, and B. F. Clarke, “Screening for thyroid dysfunction in diabetics,” *British Medical Journal*, vol. 2, no. 6202, p. 1439, 1979. View at Google Scholar · View at Scopus
 11. J. G. Hollowell, N. W. Staehling, W. Dana Flanders et al., “Serum TSH, T4, and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III),” *Journal of Clinical Endocrinology and Metabolism*, vol. 87, no. 2, pp. 489–499, 2002. View at Publisher · View at Google Scholar · View at Scopus
 12. M. F. Celani, M. E. Bonati, and N. Stucci, “Prevalence of abnormal thyrotropin concentrations measured by a sensitive assay in patients with type 2 diabetes mellitus,” *Diabetes Research*, vol. 27, no. 1, pp. 15–25, 1994. View at Google Scholar · View at Scopus
 13. T. G. A. Strieder, M. F. Prummel, J. G. P. Tijssen, E. Endert, and W. M. Wiersinga, “Risk factors for and prevalence of thyroid disorders in a cross-sectional study among healthy female relatives of patients with autoimmune thyroid disease,” *Clinical Endocrinology*, vol. 59, no. 3, pp. 396–401, 2003. View at Publisher · View at Google Scholar · View at Scopus
 14. C. Wang, “The relationship between type 2 diabetes mellitus and related thyroid diseases,” *Journal of Diabetes Research*, vol. 2013, Article ID 390534, 9 pages, 2013. View at Publisher · View at Google Scholar

15. R. W. V. Flynn, T. M. MacDonald, A. D. Morris, R. T. Jung, and G. P. Leese, "The thyroid epidemiology, audit, and research study: thyroid dysfunction in the general population," *Journal of Clinical Endocrinology and Metabolism*, vol. 89, no. 8, pp. 3879–3884, 2004. View at Publisher · View at Google Scholar · View at Scopus
16. M. A. Michalaki, A. G. Vagenakis, A. S. Leonardou et al., "Thyroid function in humans with morbid obesity," *Thyroid*, vol. 16, no. 1, pp. 73–78, 2006. View at Publisher · View at Google Scholar · View at Scopus
17. K. Boelaert, P. R. Newby, M. J. Simmonds et al., "Prevalence and relative risk of other autoimmune diseases in subjects with autoimmune thyroid disease," *American Journal of Medicine*, vol. 123, no. 2, pp. 183–e1, 2010. View at Publisher · View at Google Scholar · View at Scopus
18. R. W. Holl, B. Böhm, U. Loos, M. Grabert, E. Heinze, and J. Homoki, "Thyroid autoimmunity in children and adolescents with type 1 diabetes mellitus. Effect of age, gender and HLA type," *Hormone Research*, vol. 52, no. 3, pp. 113–118, 1999. View at Google Scholar · View at Scopus
19. G. E. Umpierrez, K. A. Latif, M. B. Murphy et al., "Thyroid dysfunction in patients with type 1 diabetes: a longitudinal study," *Diabetes Care*, vol. 26, no. 4, pp. 1181–1185, 2003. View at Publisher · View at Google Scholar · View at Scopus
20. S. A. P. Chubb, W. A. Davis, and T. M. E. Davis, "Interactions among thyroid function, insulin sensitivity, and serum lipid concentrations: the Fremantle diabetes study," *Journal of Clinical Endocrinology and Metabolism*, vol. 90, no. 9, pp. 5317–5320, 2005. View at Publisher · View at Google Scholar · View at Scopus
21. O. Kordonouri, A. Klinghammer, E. B. Lang, A. Grüters-Kieslich, M. Grabert, and R. W. Holl, "Thyroid autoimmunity in children and adolescents with type 1 diabetes: a multicenter survey," *Diabetes Care*, vol. 25, no. 8, pp. 1346–1350, 2002. View at Publisher · View at Google Scholar · View at Scopus
22. J.-K. Yang, W. Liu, J. Shi, and Y.-B. Li, "An association between subclinical hypothyroidism and sight-threatening diabetic retinopathy in type 2 diabetic patients," *Diabetes Care*, vol. 33, no. 5, pp. 1018–1020, 2010. View at Publisher · View at Google Scholar · View at Scopus
23. M. P. J. Vanderpump, W. M. G. Tunbridge, J. M. French et al., "The incidence of thyroid disorders in the community: a twenty-year follow-up of the Wickham Survey," *Clinical Endocrinology*, vol. 43, no. 1, pp. 55–68, 1995. View at Google Scholar · View at Scopus
24. N. Friedrich, S. Schwarz, J. Thonack, U. John, H. Wallaschofski, and H. Völzke, "Association between parity and autoimmune thyroiditis in a general female population," *Autoimmunity*, vol. 41, no. 2, pp. 174–180, 2008. View at Publisher · View at Google Scholar · View at Scopus
25. F. Santini, L. Chiovato, R. Rocchi, M. Marinò, and A. Pinchera, "Influences of thyroid diseases in diabetic pregnant women," *Annali dell'Istituto Superiore di Sanita*, vol. 33, no. 3, pp. 441–445, 1997. View at Google Scholar · View at Scopus
26. IDF Diabetes Atlas 5th edition 2012 update," International Diabetes

Federation,<http://www.idf.org/diabetesatlas/5e/the-global-burden>.

27. American diabetes Association, “Standards of Medical care in diabetes—2010,” *Diabetes Care*, vol. 33, supplement 1, pp. S11–S61, 2010. View at Publisher · View at Google Scholar
28. G. Caldwell, H. A. Kellett, and S. M. Gow, “A new strategy for thyroid function testing,” *The Lancet*, vol. 1, no. 8438, pp. 1117–1119, 1985. View at Google Scholar · View at Scopus
29. H. Lindner, “The Non-Clinical ATA/AACE 2012 Hypothyroidism Guidelines,”<http://hormonerestoration.com/files/ResponsetoGuidelines.pdf>.